

Häst Brunet Garde

ID 98124

Sid 2/3

Journal

2023-06-01 Surgery: dorsal recumbence. Arthroscopy of the LH tibiotarsal joint (TTJ) and LH metatarsophalangeal joint (MTJ).
 Operation
 KaBa LH TTJ- OCD at the level of the distal third of the lateral talus ridge, 1 fragment circa 4 x 6 mm large.
 Rest of the joint is in a good condition. No signs of OA.
 LH MTJ:
 - POF at the level of the plantaroproximal and medial aspect of the P1, 1 fragment circa 5 mm long.
 Rest of the joint is in a good condition. No signs of OA.
 - non- united epicondyle with intraarticular component visible at the level of the plantaroproximal and lateral aspect of the P1. This fragment is minimally unstable with no signs of intraarticular damage to the ipsilateral (lateral) condyle of the MTT III. This fragment should be radiographically monitored and the horse eventually undergoes osteosynthesis.

The fragments from the LH MTJ and LH TTJ were removed and the OCD bed debrided with sharp curettage and shaver. Joints were thoroughly lavaged with sterile saline. Arthroscopic portals were sutured with Monocryl 2-0USP/3M using simple interrupted suture pattern. High and distal limb bandages were placed on postoperatively. Uneventful spontaneous recovery from the general anesthesia.

NSAID (Rheumocam) orally for 5-7 days. Bandage change every 5-7 days (based on how the bandage looks like) for a total period of 3 weeks post op.
 Suture removal: in 2 weeks under strictly aseptic manner (following chlorhexidine or povidone iodide scrub). Afterwards place again a single layer bandage on for additional 5-7 days.
 Complete boxrest for 3 weeks (as long as the horse has a bandage on). Start with controlled handwalking exercise for 5-10 minutes per day on week 4, and add extra 5-10 minutes every other week (20 minutes on week 5, etc) to end up on 50 minutes of handwalking on week 8. Afterwards the horse can slowly resume normal exercise/be turned out into a large paddock.

2023-06-02 Morgonstatus
 Journal
 MeDe AT mycket rädd, försök att sparka
 HR 36/min, ausk cor och pulmones ua,
 Sih rosa, fuktiga, CRT<2 sek.
 GI +++++/++, avf ++
 Lnn submand och jugularer ua,
 VB bandaget sitter inte perfekt, ohalt skritt

Plan: BB och hem

2023-06-02 Bandagebyte innan hemgång. Sederas 0.3 det + 0.3 but. Lägger nytt cellonabandage över has samt kota vänster bak. Melolin 10x20 mot suturer. Suturer torra samt fina enligt MeDe.
 Journal
 FeOI

2023-06-13 Scannat finns, scan20230613_20230613093558.jpg.
 Journal

SLAKTKARENS:

Startkarens gäller i enlighet med respektive förbunds regler.

Om något oförutsett händer under icke ordinarie öppettid och som har samband med hästens behandling, ring 08-592 540 10. Vid feber eller svullnad i behandlad led bör du ringa till kliniken omgående.